



SURVEILLANCE REPORT

Bi-weekly influenza surveillance overview

26 August 2011 (erratum 14 September 2011)

Main surveillance developments in weeks 32–33/2011 (08–21 August 2011)

This first page contains the main developments during these weeks and can be printed separately or together with the more detailed information following.

- During weeks 32–33/2011, all 17 reporting countries experienced low intensity of influenza activity.
- Sentinel specimens yielded no influenza virus detections while six type A and seven type B influenza viruses were detected in non-sentinel specimens. Four of the type A viruses were subtyped, three as A(H1)2009 and one as A(H3).
- During weeks 32–33/2011, two cases of SARI were reported.

Sentinel surveillance of influenza-like illness (ILI)/ acute respiratory infection (ARI): During weeks 32–33/2011, all 17 reporting countries experienced low intensity of influenza activity. For more information, [click here](#).

Virological surveillance: Of 41 sentinel specimens, none yielded influenza detections while six type A and seven type B influenzas were detected in non-sentinel specimens. For more information, [click here](#).

Hospital surveillance of severe acute respiratory infection (SARI): Two cases of SARI were reported during weeks 32–33/2011. For more information, [click here](#).

Erratum. The following correction was made on 14 September 2011:

Page 7, first paragraph: 'In weeks 32–33/2010, 16 countries reported virological data' was changed to read: 'In weeks 32–33/2011, 16 countries reported virological data.'

Sentinel surveillance (ILI/ARI)

Weekly analysis – epidemiology


During weeks 32–33/2011, all 17 reporting countries experienced low intensity of influenza activity (Table 1, Map 1).

In terms of geographic spread, sporadic activity was reported by Cyprus and the UK (Scotland). No activity was reported by the remaining 16 countries and the UK (England and Northern Ireland).

Stable or decreasing trends were reported by all 17 countries (Table 1, Map 2).

Map 1: Intensity for weeks 32–33/ 2011***Intensity**

- [Grey square] No report
- [Yellow square] Low
- [Orange square] Medium
- [Red square] High
- [Dark red square] Very High



Liechtenstein

Luxembourg

Malta

(C) ECDC/Dundas/TESSy

* A type/subtype is reported as dominant when > 40 % of all samples are positive for the type/subtype.


Legend:

No report	Intensity level was not reported	+	Increasing clinical activity
Low	No influenza activity or influenza at baseline levels	-	Decreasing clinical activity
Medium	Usual levels of influenza activity	=	Stable clinical activity
High	Higher than usual levels of influenza activity	A(H1)2009	Type A, Subtype (H1)2009
Very high	Particularly severe levels of influenza activity		

*The map displays data for the most recent reported week only of the two-week surveillance period. For information on the other week please consult the weekly 'Influenza activity maps' [here](#).

Map 2: Geographic spread for weeks 32–33/ 2011***Geographic spread**

- [Grey square] No Report
- [Yellow square] No Activity
- [Yellow square with yellow border] Sporadic
- [Orange square] Local
- [Red square] Regional
- [Dark red square] Widespread



(C) ECDC/Dundas/TESSy

* A type/subtype is reported as dominant when at least ten samples have been detected as influenza positive in the country and of those > 40 % are positive for the type/subtype.

Legend:

No report	Activity level was not reported	+	Increasing clinical activity
No activity	No evidence of influenza virus activity (clinical activity remains at baseline levels)	-	Decreasing clinical activity
Sporadic	Isolated cases of laboratory confirmed influenza infection	=	Stable clinical activity
A(H1)2009			Type A, Subtype (H1)2009
Local outbreak	Increased influenza activity in local areas (e.g. a city) within a region, or outbreaks in two or more institutions (e.g. schools) within a region (laboratory confirmed)		
Regional activity	Influenza activity above baseline levels in one or more regions with a population comprising less than 50% of the country's total population (laboratory confirmed)		
Widespread	Influenza activity above baseline levels in one or more regions with a population comprising 50% or more of the country's population (laboratory confirmed)		

*The map displays data for the most recent reported week only of the two-week surveillance period. For information on the other week please consult the weekly 'Influenza activity maps' [here](#).

Table 1: Epidemiological and virological overview by country, weeks 32–33/ 2011*

Country	Intensity	Geographic spread	Trend	No. of sentinel swabs	Dominant type	Percentage positive*	ILI per 100 000	ARI per 100 000	Epidemiological overview	Virological overview
Austria	Low	No activity	Stable	0	None	0.0	-	-	Graphs	Graphs
Belgium	Low	No activity	Stable	-	-	0.0	21.9	662.4	Graphs	Graphs
Bulgaria	Low	No activity	Decreasing	0	None	0.0	-	228.2	Graphs	Graphs
Cyprus	Low	Sporadic	Stable	-	-	0.0	-*	-*	Graphs	Graphs
Czech Republic				-	-	0.0	-	-		
Denmark				1	None	0.0	-	-	Graphs	Graphs
Estonia	Low	No activity	Stable	1	None	0.0	1.3	64.7	Graphs	Graphs
Finland	Low	No activity	Stable	20	None	0.0	-	-	Graphs	Graphs
France				-	-	0.0	-	-		
Germany	Low	No activity	Stable	1	None	0.0	-	362.3	Graphs	Graphs
Greece	Low	No activity	Stable	-	-	0.0	14.9	-	Graphs	Graphs
Hungary				-	-	0.0	-	-		
Iceland				0	-	0.0	-	-	Graphs	Graphs
Ireland	Low	No activity	Stable	0	None	0.0	0.6	-	Graphs	Graphs
Italy				-	-	0.0	-	-		
Latvia				-	-	0.0	-	-		
Lithuania	Low	No activity	Stable	0	None	0.0	0.0	79.8	Graphs	Graphs
Luxembourg				-	-	0.0	-	-		
Malta				-	-	0.0	-	-		
Netherlands	Low	No activity	Stable	2	None	0.0	10.1	-	Graphs	Graphs
Norway	-	No activity	-	0	None	0.0	-	-	Graphs	Graphs
Poland	Low	No activity	Stable	0	None	0.0	4.3	-	Graphs	Graphs
Portugal	Low	No activity	Stable	-	-	0.0	0.0	-	Graphs	Graphs
Romania	Low	No activity	Stable	0	None	0.0	0.0	435.1	Graphs	Graphs
Slovakia	Low	No activity	Decreasing	0	A(H1)2009	0.0	32.4	471.0	Graphs	Graphs
Slovenia	Low	No activity	Stable	1	None	0.0	0.0	302.8	Graphs	Graphs
Spain				0	None	0.0	-	-	Graphs	Graphs
Sweden				0	None	0.0	-	-	Graphs	Graphs
UK - England	Low	No activity	Stable	14	None	0.0	2.5	204.4	Graphs	Graphs
UK - Northern Ireland	Low	No activity	Stable	1	-	0.0	5.3	196.3	Graphs	Graphs
UK - Scotland	Low	Sporadic	Stable	0	None	0.0	3.7	96.4	Graphs	Graphs
UK - Wales				-	-	0.0	-	-		
Europe				41		0.0			Graphs	

*Incidence per 100 000 is not calculated for these countries as no population denominator is provided.

Liechtenstein is not reporting to the European Influenza Surveillance Network.

For intensity, geographic spread and trend, the table only displays data for the most recently reported week of the two-week surveillance period.

Description of the system

Surveillance is based on nationally organised sentinel networks of physicians, mostly general practitioners (GPs), covering at least 1 to 5% of the population in their countries. All EU/EEA Member States (except Liechtenstein) participate. Depending on their country's choice, each sentinel physician reports the weekly number of patients seen with influenza-like illness (ILI), acute respiratory infection (ARI), or both to a national focal point. From the national level, both numerator and denominator data are then reported to the European Surveillance System (TESSy) database. Additional semi-quantitative indicators of intensity, geographic spread, and trend of influenza activity at the national level are also reported.

Virological surveillance

Weekly analysis – virology

In weeks 32–33/2010, 16 countries reported virological data. None of the 41 sentinel specimens tested were positive for influenza virus. Of the non-sentinel source specimens, e.g. specimens collected for diagnostic purposes in hospitals, 13 were positive for influenza virus, six were type A and seven were type B. Four type A influenza viruses were subtyped, three as A(H1)2009 and one as A(H3).

Of the 58 929 influenza viruses detected since week 40/2010 in sentinel and non-sentinel specimens, 38 966 (66.1%) were influenza A and 19 963 (33.9%) were influenza B viruses. Of 28 016 influenza A viruses subtyped, 27 236 (97.2%) were A(H1)2009 and 780 (2.8%) were A(H3) viruses (Table 2).

Since week 40/2010, 4 707 influenza viruses from sentinel and non-sentinel specimens have been characterised antigenically (Figure 1): 2 350 as A/California/7/2009 (H1N1)-like; 2 010 as B/Brisbane/60/2008-like (Victoria lineage), 185 as B/Florida/4/2006-like (Yamagata lineage), 149 as A/Perth/16/2009 (H3N2)-like and 13 as B/Bangladesh/3333/2007-like (Yamagata lineage).

More details on circulating viruses can be found in the [July](#) report prepared by the Community Network of Reference Laboratories (CNRL) coordination team. The viruses circulating this season remain well-matched with the 2010/11 seasonal vaccine viruses.


Based on centrally generated antiviral testing results now submitted to TESSy, some data for an expanded number of countries has become available. Overall, in terms of testing for oseltamivir resistance, 2562 A(H1)2009, 100 A(H3) and 508 influenza B viruses have been screened. Of these, 117 A(H1)2009 viruses were resistant with the bulk being detected in the UK (58), Denmark (27) and France (12). All viruses tested remained sensitive to zanamivir. In terms of adamantane sensitivity, all A(H1)2009 and A(H3) viruses (261 and 53 screened, respectively) were resistant.

Table 2: Weekly and cumulative influenza virus detections by type, subtype and surveillance system, weeks 40/2010–33/2011

Virus type/subtype	Current Period		Season		
	Sentinel	Non-sentinel	Sentinel	Non-sentinel	
Influenza A	0	6	8445	30521	
A (H1) 2009	0	3	7591	19645	
A (subtyping not performed)	0	2	593	10357	
A (not subtypable)	0	0	0	0	
A (H3)	0	1	261	519	
A (H1)	0	0	0	0	
Influenza B	0	7	5741	14222	
Total Influenza	0	13	14186	44743	

Note: A(H1)2009, A(H3) and A(H1) includes both N-subtyped and non-N-subtyped viruses

Figure 1: Results of antigenic characterisations of sentinel and non-sentinel influenza virus isolates, weeks 40/2010–33/2011



Description of the system

According to the nationally defined sampling strategy, sentinel physicians take nasal or pharyngeal swabs from patients with influenza-like illness (ILI), acute respiratory infection (ARI) or both and send the specimens to influenza-specific reference laboratories for virus detection, (sub-)typing, antigenic or genetic characterisation and antiviral susceptibility testing.

For details on the current virus strains recommended by WHO for vaccine preparation [click here](#).

Hospital surveillance – severe acute respiratory infection (SARI)

Weekly analysis – SARI

During weeks 32–33/2011, two cases of SARI were reported.

Since week 40/2010, 5 238 SARI cases including 513 related fatalities have been reported by 10 countries (Table 3). Of the 3 843 hospitalised cases with confirmed influenza virus infection, 3 527 (91.8%) were type A and 316 (8.2%) were type B. Of the 3 119 subtyped influenza A viruses, 3 096 (99.3%) were A(H1)2009 and 23 (0.7%) were A(H3) (Table 4). Since week 40/2010, of 2 062 SARI cases admitted to ICU, at least 1 042 (50.5%) required ventilation (Table 5).

Table 3: Cumulative number of SARI cases, weeks 40/2010–33/2011

Country	Number of cases	Incidence of SARI cases per 100 000 population	Number of fatal cases reported	Incidence of fatal cases per 100 000 population	Estimated population covered
Finland	94		30		
Romania	449	7	30	0.47	6413821
Austria	505		26		
France	790		144		
Ireland	122		23		
Portugal	418		45		
Belgium	952				
Spain	1621		192		
Malta	55	13.3	1	0.24	413609
Slovakia	232	4.27	22	0.4	5435273
Total	5238		513		

Table 4: Number of SARI cases by influenza type and subtype, weeks 30-33/ 2011 and cumulative for the season

Virus type/subtype	Number of cases during current week	Cumulative number of cases since the start of the season
Influenza A		3527
A(H1)2009		3096
A(subtyping not performed)		408
A(H3)		23
Influenza B		316
Other Pathogen		39
Unknown	2	1356
Total	2	5238

Table 5: Number of SARI cases by level of care and respiratory support, weeks 40/2010–33/2011

Respiratory support	ICU	Inpatient ward	Other	Unknown
No respiratory support necessary	178	573	449	
Oxygen therapy	144	205	395	
Respiratory support given unknown	698	339	882	251
Ventilator	1042	22	6	54

This report was written by an editorial team at the European Centre for Disease Prevention and Control (ECDC): Eeva Broberg, Flaviu Plata and René Snacken. The bulletin text was reviewed by the Community Network of Reference Laboratories for Human Influenza in Europe (CNRL) coordination team: Adam Meijer, Rod Daniels, John McCauley and Maria Zambon. On behalf of the EISN members, the bulletin text was reviewed by Bianca Snijders (RIVM Bilthoven, Netherlands) and Thedi Ziegler (National Institute for Health and Welfare, Finland). In addition, the report is reviewed by experts of WHO Regional Office for Europe.

Maps and commentary published in this Weekly Influenza Surveillance Overview (WISO) do not represent a statement on the part of ECDC or its partners on the legal or border status of the countries and territories shown.

All data published in the WISO are up-to-date on the day of publication. However, after this date published data should not be used for longitudinal comparisons as countries tend to retrospectively update their databases.

© European Centre for Disease Prevention and Control, Stockholm, 2011