

ECDC TECHNICAL REPORT

Infection prevention and control measures for Ebola virus disease Management of healthcare workers returning from Ebola-affected areas

21 January 2015

Proposed options

All healthcare workers returning from affected areas should be provided with information upon their return. In addition they should undergo an individual exposure assessment as early as possible upon returning. Additional measures can be considered on the basis of the results of the exposure assessment, using the guidance in the table below.

Type of exposure	Proposed option(s) for measures
No direct contact with Ebola patients or their bodily fluids (e.g. involved in training local healthcare workers)	Passive monitoring
Appropriately protected contact with bodily fluids of Ebola patients (e.g. laboratory worker), fomites (e.g. bed linen) or during clinical activities	Active monitoring
Unprotected, inappropriately protected contact or known breach of protection while caring for an Ebola patient, handling bodily fluids of a patient or fomites	Active monitoring Restriction of engagement in clinical activities No travel abroad
Mucosa or parenteral direct contact with bodily fluids of an Ebola patient (e.g. pricking a finger with a needle used for a patient or getting bodily fluid projection in the eyes)	Active monitoring Restriction of engagement in clinical activities Restriction of social interactions Restriction of movement

The proposed options in the table above should be modulated according to the individual risk assessment. Healthcare workers need to be informed prior to departure of the measures they should expect upon returning from affected areas. Upon returning from affected areas, all healthcare workers should be informed of the procedures for their monitoring, following an individual assessment of exposures.

Measures for returning HCWs should be proportionate to the risks, and ensure the best possible management in case they become ill, while aiming to protect public health and in respect of their personal rights. Current evidence on transmission, particularly that it has only been clearly documented from symptomatic patients, suggests that full quarantine of asymptomatic contacts, including healthcare workers, would be a disproportionate measure, unless there is evidence of non-compliance with recommended measures resulting in a potential risk to public health.

Background

The unprecedented magnitude and geographic extent of the Ebola virus disease (EVD) epidemic in West Africa has overwhelmed the local response capacity, posing an extreme challenge for epidemic containment [1]. This has led to an international response which has increased over time, although the epidemic is not as yet contained.

The increasing international response has meant that a large number of European and international experts, particularly healthcare workers, have been deployed in affected countries through the Global Outbreak Alert and Response Network (GOARN) and various governmental and non-governmental organisations. Médecins Sans Frontières (MSF) has deployed over 700 individuals in the field [2] and many countries have deployed medical and military staff to the region [3-5].

Healthcare workers have been identified as being particularly at risk for Ebola infection and 843 have been infected, with 500 deaths in affected West African countries as of 11 January 2015 [6]. The majority of these were local healthcare workers and many of these infections occurred following exposures in non-specialised units or in the community, rather than in Ebola treatment centres. A number of international healthcare workers have also been infected, with 12 having been medically evacuated to Europe for treatment, as of 16 January 2015 [7].

Healthcare workers are at risk of infection at various points in healthcare settings:

- in the early stage of symptomatic disease when unprotected contact can occur, as patients are not very contagious [8] but their infection is not yet recognised
- in the later stages of disease, after EVD is confirmed, when patients may experience very high viral loads while undergoing contamination-prone invasive procedures as part of their medical care. This has been the case in Spain and the United States while taking care of repatriated or imported EVD cases [9,10].

Following the declaration of the Public Health Event of International Concern (PHEIC) on 8 August 2014, WHO recommended that affected countries conduct exit screening of all persons at international airports, seaports and major land crossings for unexplained febrile illness consistent with potential Ebola infection. WHO also recommended that there should be no international travel of known Ebola cases or contacts of cases, unless the travel is part of an appropriate medical evacuation [11]. Although these recommendations do not specifically address healthcare workers, any symptomatic healthcare workers are unlikely to travel outside of a medical evacuation. However, due to the 21-day maximum incubation period of EVD [8], healthcare workers might travel and develop symptoms in the days after leaving affected areas as happened in the USA, where a healthcare worker was screened on arrival at a New York airport and was asymptomatic, but developed symptoms days after arrival [12].

This document outlines current monitoring processes for returning healthcare workers and provides different options for reducing the risk of transmission of EVD from infected healthcare workers returning to the EU/EEA.

Existing guidance

Public health agencies and non-governmental organisations have issued recommendations on measures which should be taken by healthcare workers upon returning from Ebola-affected areas. The following section summarises guidance issued by selected agencies and organisations.

US Centers for Disease Control and Prevention (CDC)

The Centers for Disease Control and Prevention (CDC) in the United States (USA) issued interim guidance for monitoring and movement of persons potentially exposed to Ebola virus [13]. The guidelines are based on risk categories. Any person who has unprotected exposure to blood or bodily fluids of a symptomatic case without personal protective equipment (PPE) is considered as 'high risk'. Healthcare workers using appropriate PPE in countries with widespread transmission are considered as having 'some risk', whereas healthcare workers using appropriate PPE outside countries with widespread transmission are considered as having 'low but not zero risk'.

Healthcare workers caring for symptomatic Ebola patients while wearing appropriate PPE (irrespective of whether this was in West Africa or the USA) are required to undergo 'direct active monitoring' which means that an official from the relevant public health authority assesses daily the presence of symptoms and fever through direct observation for 21 days following the return from an affected area.

Healthcare workers in the 'low but not zero risk' category do not have any restrictions imposed on travel, work, attendance at public gatherings or public transport use. Those in the 'some risk' category are assessed individually and might have restrictions related to the use of public transport, long-distance travel (airplane, bus, ship, train), attendance at mass gatherings or public places and workplaces. Persons in the 'high risk' category are restricted from the use of public transport, long-distance travel, attendance at mass gatherings or public places and workplaces.

Following the identification of the case in a returning healthcare worker in New York, the states of Florida, Illinois, New Jersey and New York issued a mandatory 21-day quarantine on healthcare workers returning from affected areas [14-16]. Mandatory quarantine of healthcare workers has been criticised by the New York Health Commissioner and by other experts as it might lead to difficulties recruiting staff to work in affected areas [17,18] besides affecting their personal rights [19].

Public Health Agency of Canada

The Public Health Agency of Canada recommends that returning healthcare workers who have used appropriate PPE and not suspected any breach should self-monitor for symptoms of Ebola for 21 days from the last contact with an Ebola case [20]. Symptomatic persons should contact their doctor prior to a planned visit to allow for safe management. Returning healthcare workers who 'provided care to cases or who have had other close physical contact with the case or deceased body, that may have resulted in unprotected exposure to blood or other body fluids from the case' are required to check their temperature twice a day and report a temperature above 38.6 °C, refrain from taking antipyretic medication, self-monitor for symptoms and self-isolate in case of any symptoms. They are advised that they should contact public health services as quickly as possible if any symptoms develop. No work restrictions are mentioned for asymptomatic healthcare workers.

Public Health England

Public Health England (PHE), on 9 January 2015, updated the recommendation for workers returning from affected areas on the degree of follow-up and work activities which may be undertaken in the following 21 days [21]:

- Persons who did not have contact with an Ebola case or bodily fluids from a case are allowed to return to normal activities without the need for monitoring or any reporting requirements.
- Persons who have been in direct contact with Ebola cases and/or bodily fluids from cases, but with use of appropriate PPE and without knowledge of any breaches, are allowed to return to clinical work, but should avoid exposure-prone procedures. They are also required to check their temperature for 21 days following return and to report any temperature over 37.5 °C.
- Persons who had direct contact with Ebola cases and/or bodily fluids from cases, but who suspect having had breaches in PPE and/or did not use appropriate PPE, are allowed to return to normal activities with some restrictions: They may use only private accommodation and, after returning from the airport, can take local public transport for up to two hours and travel by private car for an unlimited duration of time. Air travel during this period is not acceptable. They are required to check their temperature twice daily for 21 days after return, and to report daily to a designated person, even if they do not have a raised temperature (over 37.5 °C) or other suspicious symptoms. Health care workers are not allowed any patient contact for the period of 21 days.
- Symptomatic persons should call a specific contact number or the National Health Service or emergency services and provide information about symptoms and travel history to allow for safe management.

Médecins Sans Frontières (MSF)

Médecins Sans Frontières, as one of the major organisations deploying healthcare workers in West Africa, has issued recommendations on self-monitoring for its workers [2]. The recommendations for the 21 days following return from affected areas include:

- checking temperature twice a day
- finishing a regular course of malaria prophylaxis (malaria symptoms can mimic Ebola symptoms)
- being aware of relevant symptoms, such as fever
- staying within four hours of a hospital with isolation facilities
- immediately contacting MSF offices if any relevant symptoms develop.

MSF does not recommend self-quarantine of returning staff. However, returned staff members are discouraged from returning to work during the 21-day period and MSF continues to provide salaries to returned staff during this time.

ECDC

The ECDC contact management document considers follow-up of healthcare workers who have cared for EVD cases as a specific group due to the continuous nature of occupational exposure even when using appropriate PPE [22].

Healthcare workers who have cared for EVD cases in EU/EEA hospitals should be registered and monitored as part of the occupational health practices of their country of practice. This usually involves registration, active monitoring of symptoms and a prompt investigation in case of any symptoms possibly related to EVD.

Potential measures for management of healthcare workers returning from affected areas

This section summarises the potential measures that can be considered for the management of asymptomatic healthcare workers returning from affected areas.

Registration

Establishing a register of healthcare workers engaged in providing care for EVD patients in affected countries and areas can facilitate contact with them and potential monitoring of their status on return.

Information to returning healthcare workers

The information provided should include:

- general information on EVD: incubation period, clinical presentation, transmission
- advice on general protective measures for contacts, with specific attention to family and close friends as well as co-workers
- advice on the monitoring regime recommended after deployment
- procedures and information on how to report symptoms and seek medical help
- a contact number for the responsible public health office, available 24 hours a day, in case of onset of symptoms.

Healthcare workers should be advised to call their doctor or hospital should they develop symptoms within the 21-day period following their departure from an affected area prior to attending, and inform them of their travel history and possible exposure.

Individual exposure assessment

This measure consists of conducting an individual assessment of healthcare workers returning from affected areas, taking into account:

- the nature of activities in affected areas: clinical, laboratory, epidemiological
- possible exposure to EVD cases or bodily fluids using recommended personal protection procedures and equipment
- possible unprotected exposures having taking place
- known breach of personal protective measures.

The assessment should characterise the level of exposure for the healthcare worker and should be used as the basis for determining the measures to be applied.

The assessment should result in the categorisation of the healthcare worker exposure, in order of magnitude of risk:

- no direct contact with EVD patients or their bodily fluids (e.g. involved in training local healthcare workers)
- appropriately protected contact with bodily fluids of EVD patients (e.g. laboratory worker), fomites (e.g. bed linen) or during clinical activities
- unprotected, inappropriately protected contact or known breach of protection while caring for an EVD patient, handling bodily fluids of a patient or fomites
- mucosa or parenteral direct contact with bodily fluids of a patient (e.g. pricking a finger with a needle used for a patient or getting bodily fluid projection in the eyes).

The assessment provides an opportunity to offer psychological support to returning healthcare workers.

Monitoring of symptoms and body temperature

This measure relies on monitoring of symptoms and body temperature twice a day for 21 days after the last possible exposure. Monitoring can be:

- **passive:** self-monitoring by returning healthcare worker
- **active:** the returning healthcare worker having to report daily the result of the monitoring to a health authority or to the employer
- **direct active:** monitoring is done through direct observation of the healthcare worker by a health officer.

Monitoring can use forms and checklists to ensure consistency with different healthcare workers over time.

In case of fever of any grade or other symptoms compatible with EVD, the healthcare worker should immediately self-isolate and contact health services.

Restriction of engagement in clinical activities

Clinical activities are situations where transmission of disease can occur from a healthcare worker to a patient, because of the nature of the care provided and the status of the patient. This measure ensures that a healthcare worker does not perform activities that will involve contact with patients.

Restriction of social interactions

This measure imposes voluntary social interaction limitations. It may be complemented by the maintenance by the healthcare worker of a register of contact occurring during the 21-day monitoring phase. It may require non-attendance at the workplace during the monitoring period.

Restriction of movement

This measure considers the limitation of use of public transport and attendance at public events and celebrations. It may require that no travel abroad takes place or that the healthcare worker remains within four hours of a health facility with isolation capacity.

Quarantine (self/mandatory)

This measure requires that the returning healthcare worker remains confined at home (self-quarantine) or in a dedicated facility (mandatory quarantine) for the duration of the monitoring. This measure results in a minimal number of interactions with contacts. This measure should be accompanied by psychosocial support and financial compensation should be considered.

Scientific evidence

The scientific evidence underlining the effectiveness of the measures indicated in the previous section is limited. Scientific evidence on the use of temperature for screening for infection is referenced in the ECDC technical document on 'Exit and entry screening measures' [25]. Scientific evidence regarding the characteristics of the disease and its mode of transmission is referenced in the ECDC rapid risk assessments [1]. Measures related to restriction of movements and social interactions, including quarantine are effective during the period of infectiousness of a patient, but there is no scientific evidence supporting their appropriateness in the asymptomatic phase of the disease.

Proposed options

All healthcare workers returning from affected areas should be provided with information upon their return. In addition they should undergo an individual exposure assessment as early as possible upon returning. Additional measures can be considered on the basis of the results of the exposure assessment, using the guidance in the table below.

Type of exposure	Proposed option(s) for measures
No direct contact with Ebola patients or their bodily fluids (e.g. involved in training local healthcare worker)	Passive monitoring
Appropriately protected contact with bodily fluids of Ebola patients (e.g. laboratory worker), fomites (e.g. bed linen) or during clinical activities	Active monitoring
Unprotected, inappropriately protected contact or known breach of protection while caring for an Ebola patient, handling bodily fluids of a patient or fomites	Active monitoring Restriction of engagement in clinical activities No travel abroad
Mucosa or parenteral direct contact with bodily fluids of an Ebola patient (e.g. pricking a finger with a needle used for a patient or getting bodily fluid projection in the eyes).	Active monitoring Restriction of engagement in clinical activities Restriction of social interactions Restriction of movement Post-exposure prophylaxis can be considered

The proposed options in the table above should be modulated according to the individual risk assessment. Current evidence on transmission, particularly that it has only been clearly documented in symptomatic patients, suggests that full quarantine of asymptomatic contacts, including healthcare workers, would be a disproportionate measure, unless there is evidence of non-compliance with recommended measures resulting in a potential risk to public health. Measures adopted by Member States may also vary according to national laws and regulations.

Conclusion

Healthcare workers need to be informed prior to departure of the measures they should expect upon returning from affected areas. Upon returning from affected areas, all healthcare workers should be informed of the procedures for their monitoring, following an individual assessment of exposure. Measures for returning healthcare workers should be proportionate to the risks, ensure the best possible management in case they become ill, while aiming to protect public health and in respect of their personal rights.

References

1. European Centre for Disease Prevention and Control. Outbreak of Ebola virus disease in West Africa - rapid risk assessment. Fifth update, 29 September 2014 [Internet]. Stockholm: ECDC; 2014. Available from: <http://ecdc.europa.eu/en/publications/Publications/Ebola-Sierra%20Leone-Liberia-Guinea-Nigeria-23-09-2014-rapid-risk-assessment.pdf>.
2. Médecins Sans Frontières. MSF protocols for staff returning from Ebola-affected countries [Internet]. MSF USA; 2014 Oct 23 [cited 2014 Oct 24]. Available from: <http://www.doctorswithoutborders.org/article/msf-protocols-staff-returning-ebola-affected-countries>.
3. African Union. Over 2000 German volunteers offer to deploy to fight Ebola virus disease [Internet]. Addis Ababa: African Union; 2014 Sep 26 [cited 2014 Oct 27]. Available from: <http://cpauc.au.int/en/content/over-2000-german-volunteers-offer-deploy-fight-ebola-virus-disease>.
4. Pineau E, Samb S. U.N. to deploy Ebola mission as death toll reaches 2,630 [Internet]. Paris: Reuters; 2014 Sep 19 [cited 2014 Oct 27]. Available from: <http://uk.reuters.com/article/2014/09/18/uk-health-ebola-toll-idUKKBN0HD1I720140918>.
5. Department for International Development, Foreign & Commonwealth Office, The Rt Hon Justine Greening MP and The Rt Hon Philip Hammond MP. The UK is leading the international drive against Ebola in Sierra Leone [press release] [Internet]. London: UK government; 2014 [cited 2014 Oct 27]. Available from: <https://www.gov.uk/government/news/the-uk-is-leading-the-international-drive-against-ebola-in-sierra-leone>.
6. World Health Organization. Ebola response roadmap situation report. 22 October 2014 [Internet]. Geneva: WHO; 2014. Available from: http://apps.who.int/iris/bitstream/10665/137091/1/roadmapsitre22Oct2014_eng.pdf?ua=1.
7. European Centre for Disease Prevention and Control. Epidemiological update: Outbreak of Ebola virus disease in West Africa 2015 [updated 16 January]. Available from: http://www.ecdc.europa.eu/en/press/news/layouts/forms/News_DispForm.aspx?List=8db7286c-fe2d-476c-9133-18ff4cb1b568&ID=1151.
8. Bannister B. Viral haemorrhagic fevers imported into non-endemic countries: risk assessment and management. *Br Med Bull*. 2010;95:193-225.
9. Ministerio de Sanidad Servicios Sociales e Igualdad. Diagnosticado un caso secundario de contagio por virus Ébola [press release in Spanish] [Internet]. 2014 [cited 2014 Oct 6]. Available from: <http://www.msssi.gob.es/gabinete/notasPrensa.do?id=3427>.
10. Centers for Disease Control and Prevention. Texas reports positive test for Ebola in one additional healthcare worker [Internet]. Atlanta: CDC; 2014 Oct 15 [cited 2014 Oct 16]. Available from: <http://www.cdc.gov/media/releases/2014/s1015-texas-second-health-care-worker.html>.
11. World Health Organization. 2014 Ebola Virus Disease (EVD) outbreak in West Africa - Travel and transport risk assessment: Recommendations for public health authorities and transport sector [Internet]. Geneva: WHO; 2014 [cited 2014 29 July 2014]. Available from: <http://www.who.int/ith/updates/20140421/en/>.
12. New York City Department of Health and Mental Hygiene. Ebola Patient Timeline [Internet]. New York City: NYC Health; 2014 [cited 2014 Oct 27]. Available from: <http://www.nyc.gov/html/doh/downloads/pdf/cd/ebola-patient-timeline.pdf>.

13. Centers for Disease Control and Prevention. Interim U.S. guidance for monitoring and movement of persons with potential Ebola virus exposure [Internet]. Atlanta: CDC; 2014 Oct 27. Available from: <http://www.cdc.gov/vhf/ebola/pdf/monitoring-and-movement.pdf>.
14. New York City Governor. Governor Andrew Cuomo and Governor Chris Christie announce additional screening protocols for Ebola at JFK and Newark Liberty International Airports [Internet]. Albany, NY: Office of the New York Governor; 2014 Oct 24 [cited 2014 Oct 27]. Available from: <http://www.governor.ny.gov/press/10242014-ebola-screening-protocols>.
15. Illinois Governor. Governor Quinn directs Illinois Department of Public Health to require quarantine to protect against Ebola [Internet]. Chicago: Office of the Illinois Governor; 2014 Oct 24 [cited 2014 Oct 27]. Available from: <http://www3.illinois.gov/PressReleases/ShowPressRelease.cfm?SubjectID=2&RecNum=12783>.
16. Florida Governor. Gov. Scott Calls on feds to require same monitoring for volunteer workers as US Military to protect citizens, after new Ebola case [Internet]. Tallahassee: Office of the Florida Governor; 2014 Oct 24 [cited 2014 Oct 27]. Available from: <http://www.flgov.com/2014/10/24/gov-scott-calls-on-feds-to-require-same-monitoring-for-volunteer-workers-as-us-military-to-protect-citizens-after-new-ebola-case/>.
17. Sanchez R, Cohen E. N.Y. official: New Ebola policy 'a real stunner' [Internet]. CNN; 2014 Oct 26 [cited 2014 Oct 27]. Available from: <http://edition.cnn.com/2014/10/25/health/us-ebola/index.html>.
18. CBS/AP. Will mandatory Ebola quarantines make things worse? [Internet]. CBS; 2014 Oct 26 [cited 2014 Oct 27]. Available from: <http://www.cbsnews.com/news/ebola-outbreak-will-mandatory-health-worker-quarantines-make-things-worse/>.
19. Médecins Sans Frontières. MSF nurse held in isolation in New Jersey [Internet]. MSF; 2014 Oct 25 [cited 2014 Oct 27]. Available from: <http://www.doctorswithoutborders.org/article/msf-nurse-held-isolation-new-jersey>.
20. Public Health Agency of Canada. Interim guidance - Ebola virus disease: Infection prevention and control measures for borders, healthcare settings and self-monitoring at home [Internet]. Ottawa: Public Health Agency of Canada; 2014 [updated 2014 Sep 12; cited 2014 Oct 27]. Available from: <http://www.phac-aspc.gc.ca/id-mi/vhf-fvh/ebola-ipc-pci-eng.php#tbl-3>.
21. Public Health England. Public health recommendations for returning workers who are asymptomatic contacts of Ebola virus disease 2015. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/394131/20150109_Ebola_recommendations_for_returning_workers_who_are_asymptomatic_contacts.pdf.
22. European Centre for Disease Prevention and Control. Public health management of persons having had contact with Ebola virus disease cases in the EU [Internet]. Stockholm: ECDC; 2014 Oct 22. Available from: <http://ecdc.europa.eu/en/publications/Publications/ebola-public-health-contact-management.pdf>.